

A LOWER BOUND ON SESHADRI CONSTANTS OF HYPERPLANE BUNDLES ON THREEFOLDS

KUNGHO CHAN

ABSTRACT. We give the lower bound on Seshadri constants for the case of very ample line bundles on threefolds. We consider the situation when the Seshadri constant is strictly less than 2 and give a version of Bauer's theorem [Bau99, Theorem 2.1] for singular surfaces so we can prove the same result for smooth threefolds.

1. INTRODUCTION

The Seshadri constant at a given point on a smooth projective variety was introduced by Demailly [Dem92] to study Fujita's conjecture. It measures how positive a nef line bundle locally is near a given point. Since then Seshadri constants were recognized as interesting invariants of algebraic varieties on their own.

Definition 1.1. Suppose X is a projective variety of dimension n and L is a nef line bundle over X . Let x be a point on X and

$$\pi : \tilde{X} \rightarrow X$$

the blowup of X at x with the exceptional divisor E . Then, we define the *Seshadri constant* $\epsilon(L, x)$ of L at x as

$$\epsilon(L, x) := \sup\{\alpha \geq 0 \mid \pi^*L - \alpha E \text{ is nef}\}.$$

Or, equivalently, it can be defined by

$$\epsilon(L, x) := \inf_{x \in C \subset X} \left\{ \frac{L \cdot C}{\text{mult}_x C} \right\}$$

where the infimum is taken over all integral curves $C \subset X$ passing through x [Laz04, 5.1.5].

For the case that $X \subset \mathbf{P}^N$ is a smooth integral projective variety and L is the restriction $\mathcal{O}_X(1)$ on X of the hyperplane bundle of $\mathbf{P}^N$, it is easy to see that $\epsilon(\mathcal{O}_X(1), x) \geq 1$ for every $x \in X$. Obviously, the equality holds if there is a line in X passing through x . For the case of smooth surfaces, Bauer proved the following.

Theorem 1.2. [Bau99, Theorem 2.1] *Let $X \subset \mathbf{P}^N$ be a smooth irreducible surface.*

- (a) $\epsilon(\mathcal{O}_X(1), x) = 1$ if and only if X contains a line passing through x .
- (b) Suppose X is of degree d (≥ 4) and x is a point on X . If X contains no line passing through x , then

$$\epsilon(\mathcal{O}_X(1), x) \geq \frac{d}{d-1}.$$

Date: Received: date / Revised version: date.

- (c) If X is of degree d (≥ 4) and $x \in X$ is a point such that the Seshadri constant $\epsilon(\mathcal{O}_X(1), x)$ satisfies the inequalities $1 < \epsilon(\mathcal{O}_X(1), x) < 2$, then it is of the form

$$\epsilon(\mathcal{O}_X(1), x) = \frac{a}{b},$$

where a, b are integers with $3 \leq a \leq d$ and $a/2 < b < a$.

- (d) All rational numbers a/b with $3 \leq a \leq d$ and $a/2 < b < a$ occur as local Seshadri constants of smooth irreducible surfaces in $\mathbf{P}^3$ of degree d .

Bauer's approach was to consider the intersection of X and the tangent plane $T_x X$ of X at x . If the Seshadri constant $\epsilon(\mathcal{O}_X(1), x) < 2$, then it must be computed by a component of the intersection and the multiplicity of the component at x is bounded above by the degree d of X . In this paper, we prove a similar result for any integral surfaces in $\mathbf{P}^3$.

Proposition 1.3. *Let X be a projective integral surface of degree d (≥ 3) in $\mathbf{P}^3$. If $x \in X$ is a point of multiplicity m and X contains no line passing through x , then*

$$\epsilon(\mathcal{O}_X(1), x) \geq \frac{d}{d-1}.$$

- (a) If $\epsilon(\mathcal{O}_X(1), x) < \frac{m+1}{m}$, then

$$\epsilon(\mathcal{O}_X(1), x) = \frac{a}{b}$$

for some integers a, b such that $3 \leq a \leq md$ and $\frac{ma}{m+1} < b \leq \frac{a(d-1)}{d}$.

By constructing a singular surface in $\mathbf{P}^3$, we also show that the lower bound $\frac{d}{d-1}$ here is optimal. Furthermore, we give the same lower bound for smooth threefolds in projective spaces.

Theorem 1.4. *Let X be an irreducible smooth projective threefold of degree d (≥ 4) in $\mathbf{P}^N$, x a point in X and $T_x X$ the tangent linear subspace of $\mathbf{P}^N$ to X at x . If X contains no line through x , then*

- (a) If $\dim(T_x X \cap X) = 0$, then

$$\epsilon(\mathcal{O}_X(1), x) \geq 2.$$

- (b) If $\dim(T_x X \cap X) = 1$, then

$$\epsilon(\mathcal{O}_X(1), x) \geq \frac{d}{d-1}.$$

If furthermore $\epsilon(\mathcal{O}_X(1), x) < 2$, then $\epsilon(\mathcal{O}_X(1), x) = \frac{a}{b}$ for some integers a, b such that $3 \leq a \leq d$ and $\frac{a}{2} < b < a$.

- (c) If $\dim(T_x X \cap X) = 2$, then

$$\epsilon(\mathcal{O}_X(1), x) \geq \frac{d}{d-1}.$$

If furthermore $\epsilon(\mathcal{O}_X(1), x) < \frac{d-1}{d-2}$, then $\epsilon(\mathcal{O}_X(1), x) = \frac{a}{b}$ for some integers a, b such that $3 \leq a \leq (d-2)d$ and $\frac{(d-2)a}{d-1} < b \leq \frac{a(d-1)}{d}$.

Finally, we construct a threefold with finitely many singular points in $\mathbf{P}^4$ whose Seshadri constant of hyperplane bundle at a smooth point is $\frac{d}{d-1}$.

Acknowledgement. This paper is part of my PhD thesis. I thank my advisor L. Ein for getting me interested in this question and helping me with comments and suggestions.

2. SESHADRI CONSTANTS OF HYPERPLANE LINE BUNDLES ON SURFACES

In order to prove 1.3, we need some preparations.

Definition 2.1. Let X be a reduced irreducible projective variety of dimension n in $\mathbf{P}^N$ and x a point on X . Denote $\mathcal{O}_X(1) = \mathcal{O}_{\mathbf{P}^N}(1)|_X$ by H , the blowup of $\mathbf{P}^N$ at x by $\widetilde{\mathbf{P}}^N$, the proper transform of X in $\widetilde{\mathbf{P}}^N$ by $\widetilde{X}$ and the exceptional divisor in $\widetilde{X}$ by E . The projection $\mathbf{P}^N \setminus \{x\} \rightarrow \mathbf{P}^{N-1}$ induces a morphism $\widetilde{\mathbf{P}}^N \rightarrow \mathbf{P}^{N-1}$ such that the proper transform of every line in $\mathbf{P}^N$ passing through x is mapped to a point in $\mathbf{P}^{N-1}$. We call the restriction on $\widetilde{X}$ of this morphism the *inner projection* of $X \subset \mathbf{P}^N$ from x , that is

$$p : \widetilde{X} \rightarrow \mathbf{P}^{N-1}.$$

If $\pi : \widetilde{X} \rightarrow X$ is the blowup morphism of X at x , then from the construction $p^*\mathcal{O}(1) = \pi^*H - E$.

It is obvious that if there is no line in X passing through x , then p is finite.

Lemma 2.2. *With the above settings, there is no line in X passing through x if and only if $\epsilon(H, x) > 1$.*

Proof. The " $\Leftarrow$ " is obvious. For " $\Rightarrow$ ", we assume the hypothesis, then p is finite. This implies that $p^*\mathcal{O}(1)$ is ample and hence $\pi^*H - E$ is ample. So $\epsilon(H, x) > 1$. $\square$

Lemma 2.3. *With the above settings, if $\epsilon(H, x) > 1$, then*

$$\deg X - r \geq \text{mult}_x X$$

where $r = \deg p(\widetilde{X})$. In particular, $\deg X - 1 \geq \text{mult}_x X$.

Proof. By 2.2, X contains no line passing through x , p is finite and hence

$$\begin{aligned} \deg X - \text{mult}_x X &= \deg X + (-1)^n E^n \\ &= (\pi^*H - E)^n \\ &= p^*\mathcal{O}(1)^n \\ (1) \quad &= \deg(p) \cdot (\mathcal{O}(1)|_{p(\widetilde{X})})^n \\ &= \deg(p) \cdot \deg p(\widetilde{X}) \\ &\geq r. \end{aligned}$$

$\square$

The following two lemmas gives us a way to bound the multiplicity of a variety at a point by its degree.

Lemma 2.4. *Suppose $T \subset \mathbf{P}^N$ is an integral cone of dimension $N - 1$ over an integral hypersurface B in $\mathbf{P}^{N-1}$ with vertex at $p \in \mathbf{P}^N$. If X is an integral subvariety in T of codimension one containing p and X contains no line passing through p , then*

$$\deg X - \deg B \geq \text{mult}_p X.$$

Proof. Since the image $p(\tilde{X}) = B$ and $\epsilon(\mathcal{O}_X(1), p) > 1$, then from (1), we have $\deg X - \text{mult}_p X = p^* \mathcal{O}(1)^{N-2} \cdot \tilde{X} = \deg(p) \cdot (\mathcal{O}(1)|_B)^{N-2} = \deg p \cdot \deg B \geq \deg B$. $\square$

Lemma 2.5. *Suppose X is a hypersurface of degree d in $\mathbf{P}^{n+1}$ and x is a point of multiplicity m on X . If X contains no line passing through x , then*

$$m \leq d - n.$$

Proof. Suppose X is defined by the homogeneous polynomial f of degree d . By choosing an appropriate coordinate system, we can assume $x = [0 : 0 : \cdots : 0 : 1]$. Then

$$f = x_{n+1}^{d-1} \cdot f_1 + x_{n+1}^{d-2} \cdot f_2 + \cdots + x_{n+1}^{d-i} \cdot f_i + \cdots + f_d$$

where f_i is a homogeneous polynomial of degree i in $x_0, x_1, \dots, x_n$. Since $\text{mult}_x X = m$, then $f_m \neq 0$ and $f_i = 0$ for $i < m$. Assume $m > d - n$, then there are at most n nonzero f_i 's with $d \geq i \geq m$. Then the intersection of the cones in $\mathbf{P}^{n+1}$ defined by nonzero f_i 's is nonempty and contains line. This implies X contains a line passing through x . This is a contradiction. $\square$

Now we come up to prove 1.3.

Proof of 1.3. Let $TC_x(X) \subset \mathbf{P}^3$ be the tangent cone of X at x . Suppose the local equation of X around x is f . By choosing an appropriate coordinate system, we can assume $x = (0, 0, 0)$ and $f = f_m + f_{\geq m+1}$ where f_i is a homogeneous polynomial of degree i . Then the local equation of $TC_x(X)$ is f_m , and hence $\deg TC_x(X) = m$ and $\text{mult}_x(TC_x(X)) = m$. Since $TC_x(X)$ is spanned by all the tangent lines to X at x and X contains no line through x , $TC_x(X)$ must intersect X properly. Denote $TC_x(X) \cap X$ by D . Then $D \in |\mathcal{O}_X(m)|$.

If $\epsilon(\mathcal{O}_X(1), x) \geq \frac{m+1}{m}$, then $\epsilon(\mathcal{O}_X(1), x) \geq \frac{d}{d-1}$ since $\frac{x+1}{x}$ is decreasing for $x > 0$ and $d - 1 \geq m$ by 2.3. Hence it reduces to assume

$$(2) \quad \epsilon(\mathcal{O}_X(1), x) < \frac{m+1}{m}.$$

This implies that there exists an integral curve $x \in C \subset X$ such that $\epsilon(\mathcal{O}_X(1), x) \leq \frac{\deg C}{\text{mult}_x C} < \frac{m+1}{m}$. We claim that C must be an irreducible component of D . To this end, let us consider the blowup of $\mathbf{P}^3$ at x with the exceptional divisor E ,

$$\pi : \widetilde{\mathbf{P}^3} \rightarrow \mathbf{P}^3.$$

Let us set $T = TC_x(X)$. Then,

$$\pi^* T = \tilde{T} + mE$$

and

$$\pi^* X = \tilde{X} + mE$$

where $\tilde{T}$ and $\tilde{X}$ are respectively the proper transforms of T and X . Since $\tilde{X}|_E = \tilde{T}|_E$, then

$$(\pi|_{\tilde{X}})^* D = \tilde{D} + \text{the common components of } (\tilde{X}|_E) \text{ and } (\tilde{T}|_E) + m \cdot \tilde{X}|_E = \tilde{D} + (m+1)(\tilde{X}|_E)$$

where $\tilde{D}$ is the proper transform of D .

Let $\tilde{C}$ be the proper transform of C .

Assume C is not an irreducible component of D , then

$$\begin{aligned} m \deg C = D \cdot C &= ((\pi|_{\tilde{X}})^* D) \cdot \tilde{C} \\ &= (\tilde{D} + (m+1)(\tilde{X}|_E)) \cdot \tilde{C} \\ &\geq (m+1) \text{mult}_x C. \end{aligned}$$

This is a contradiction and thus C is an irreducible component of D .

Suppose the tangent cone $T = \sum a_i T_i$ where T_i 's are integral cones in $\mathbf{P}^3$ with vertices at x and $\sum a_i \deg T_i = m$. Then, $C \subset (T_i \cap X)$ for some i and hence

$$\deg C \leq \deg X \cdot \deg T_i = d \deg T_i.$$

Moreover, by 2.4

$$\frac{\deg C}{\text{mult}_x C} \geq \frac{\deg C}{\deg C - \deg T_i} \geq \frac{d \deg T_i}{d \deg T_i - \deg T_i} = \frac{d}{d-1}.$$

So we have $\epsilon(\mathcal{O}_X(1), x) \geq \frac{d}{d-1}$. This is for part (a).

For part (b), if (2) is satisfied, $\epsilon(\mathcal{O}_X(1), x)$ must be computed by one of the irreducible components of D . Let C be such a component and contained in an integral cone $T_i \subset T$ then

$$\epsilon(\mathcal{O}_X(1), x) = \frac{\deg C}{\text{mult}_x C}.$$

Since X contains no line passing through x , $1 < \deg C (\leq \deg D = md)$ and $(\frac{m \deg C}{m+1} <) \text{mult}_x C \leq \frac{(d-1) \deg C}{d}$. Since $\epsilon(\mathcal{O}_X(1), x) < \frac{m+1}{m} \leq 2$, the $\deg C$ cannot equal to 2 and hence $\deg C \geq 3$. $\square$

The following example indicates that the lower bound $\frac{d}{d-1}$ is optimal.

Example 2.6. For any $d \geq 4$ and $2 \leq m \leq d-2$, there exists an integral surface $S \subset \mathbf{P}^3$ containing no line through a point $0 \in S$ such that

$$\text{mult}_0 S = m \text{ and } \epsilon(\mathcal{O}(1), 0) = \frac{d}{d-1}.$$

Construction of 2.6. Roughly speaking, we construct S by prescribing its tangent cone T , and then we have to show the irreducibility of S and it contains no line through a particular point. Moreover, we also have to show the irreducibility of the intersection $S \cap T$.

Choose a general integral plane curve B of degree m and let T be the cone over B with vertex $0 \in \mathbf{P}^3$. Let the blowup of $\mathbf{P}^3$ at 0 be

$$\pi : \widetilde{\mathbf{P}^3} \rightarrow \mathbf{P}^3$$

with the exceptional divisor E and $\tilde{T}$ the proper transform of T . Consider on $\widetilde{\mathbf{P}^3}$ the line bundle

$$L := \pi^* \mathcal{O}_{\mathbf{P}^3}(d) - (d-1)E.$$

It is clear that L is globally generated.

Let V be the image of the global section restriction morphism,

$$V := \text{Image} \left(H^0(\widetilde{\mathbf{P}^3}, L) \rightarrow H^0(\widetilde{T}, L|_{\widetilde{T}}) \right).$$

Take a general effective divisor $D \in |L|$. Then $D|_E \not\supset B = \widetilde{T}|_E$. Set $Y := \pi_* D$ and $C := Y|_T$, then

$$\deg C = dm.$$

Moreover, set $\sigma := \pi|_{\widetilde{T}} : \widetilde{T} \rightarrow T$, then

$$\sigma^*(C) = (D + (d-1)E)|_{\widetilde{T}} = \widetilde{C} + (d-1)B$$

where the $\widetilde{C}$ is the proper transform of C .

Thus,

$$\text{mult}_0(C) = \widetilde{C}.B = (\sigma^*(C) - (d-1)B).B = (d-1)m.$$

Now we show the irreducibility of C . Consider the morphism induced by $|V|$,

$$\phi : \widetilde{T} \rightarrow \mathbf{P}^{\dim|V|}.$$

Since $(L|_{\widetilde{T}})^2 = (d^2 - (d-1)^2)m > 0$, the image of ϕ cannot be 1-dimensional. Then, by the Bertini theorem for general linear sections [Laz04, 3.3.1], $\widetilde{C}$ is irreducible and so is C .

With the T , Y and C constructed above, we can construct the singular surface S .

Let $U \ni 0$ be the complement of a hyperplane in $\mathbf{P}^3$, which is affine $\mathbf{A}^3$.

By choosing an appropriate coordinate system, we can assume the defining equation of T in $\mathbf{A}^3$ is $f_m = 0$ with the vertex $(0,0,0)$ and the defining equation of Y is $f_{d-1} + f_d = 0$ where f_i is a homogeneous polynomial of degree i in the variables x_1, x_2 and x_3 .

Now, we define the surface $S|_U$ by the equation

$$f = f_m + f_{d-1} + f_d = 0.$$

Clearly,

$$\text{the tangent cone } TC_0(S) = T \text{ and } S \cap T = C.$$

Since T is irreducible, then only one irreducible component of S can pass through the point 0. If S is reducible, then $\deg C < dm$. This is impossible when we choose C with $\deg C = dm$. Thus, S is irreducible and

$$\frac{\deg C}{\text{mult}_0 C} = \frac{dm}{(d-1)m} = \frac{d}{d-1} < \frac{m+1}{m}.$$

From the proof of 1.3, C computes the Seshadri constant

$$\epsilon(\mathcal{O}(1), 0) = \frac{d}{d-1}.$$

To complete the proof, it remains to show that S contains no line through 0. Assume to the contrary that S contains a line through 0. Let the line be $\ell := \{(at, bt, ct) | t \in \mathbf{C}\}$. Then,

$$f(at, bt, ct) = 0 \quad \forall t.$$

This implies that $\ell \subset (T \cap C(f_{d-1}) \cap C(f_d))$ where $C(f_j)$ is the cone in $\mathbf{A}^3$ defined by the homogeneous polynomial f_j . However, this situation can be eliminated when we made a general choice of D above. $\square$

3. SESHADRI CONSTANTS OF HYPERPLANE BUNDLES ON SMOOTH THREEFOLDS

To extend Bauer's approach to smooth threefold X , one has to consider the intersection of the 3-dimensional tangent space $T_x X$ and X . The possible components of the intersection are points, curves and surfaces. It turns out that 1.3 can deal with the surface components.

Proof of 1.4. We start from a simple argument. Suppose $C \subset X$ is an integral curve through x . If

$$(3) \quad C \not\subset T_x X \cap X,$$

one can take a hyperplane $H \subset \mathbf{P}^N$ such that $H \supset T_x X$ and $H \not\supset C$. Then

$$\deg C = (H|_X) \cdot C \geq 2 \cdot \text{mult}_x C.$$

For (a), (3) is always true for every integral curve through x . So, $\epsilon(\mathcal{O}_X(1), x) \geq 2$.

For (b), assume $\frac{\deg C}{\text{mult}_x C} < 2$. Then C must be an irreducible component of $T_x X \cap X$. Otherwise, (3) is true for C . Now $C \subset T_x X \cap X$, then $\deg C \leq d$. By 2.3

$$\frac{\deg C}{\text{mult}_x C} \geq \frac{\deg C}{\deg C - 1} \geq \frac{d}{d-1}.$$

This induces $\epsilon(\mathcal{O}_X(1), x) \geq \frac{d}{d-1}$.

In this situation, $\epsilon(\mathcal{O}_X(1), x)$ must be computed by one of the integral curves in $T_x X \cap X$ with $2 < \deg C \leq d$ and $\frac{\deg C}{2} < \text{mult}_x C < \deg C$.

For (c), assume $\frac{\deg C}{\text{mult}_x C} < 2$. We can assume that there is an irreducible and reduced surface $S \subset T_x X \cap X$ such that $C \subset S$. If not, either (3) is true for C or C is an irreducible component of $T_x X \cap X$. For the latter situation $\deg C \leq d$, we will have $\frac{\deg C}{\text{mult}_x C} \geq \frac{d}{d-1}$ as the situation in (b).

Now S is a projective surface in $T_x X (\cong \mathbf{P}^3)$ and $\deg S \leq d$. By 1.3,

$$(4) \quad \frac{\deg C}{\text{mult}_x C} \geq \epsilon(\mathcal{O}_X(1)|_S, x) \geq \frac{d}{d-1}.$$

Combining both cases, we have $\epsilon(\mathcal{O}_X(1), x) \geq \frac{d}{d-1}$.

Assume furthermore $\epsilon(\mathcal{O}_X(1), x) < \frac{d-1}{d-2}$.

Suppose $\text{mult}_x S = m$. Since $d-2 \geq m$ from 2.5,

$$\epsilon(\mathcal{O}_X(1), x) < \frac{d-1}{d-2} \leq \frac{m+1}{m}.$$

Hence, as the part (b) of 1.3, we have

$$\epsilon(\mathcal{O}_X(1), x) = \frac{a}{b}$$

for some integers a, b such that $3 \leq a \leq (d-2)d$ and $\frac{(d-2)a}{d-1} < b \leq \frac{a(d-1)}{d}$. $\square$

This result implies that if $\epsilon(\mathcal{O}_X(1), x) < \frac{d-1}{d-2}$, then the number of possible values of $\epsilon(\mathcal{O}_X(1), x)$ is finite. Indeed, this phenomenon was generally proved for surfaces. In [Ogu02], Oguiso proved that if $(f : \mathcal{X} \rightarrow \mathcal{B}, \mathcal{L})$ is a family of polarized surfaces of

degree d and $\Sigma := \{\epsilon(\mathcal{L}_t, x_t) \mid t \in \mathcal{B}, x_t \in \mathcal{X}_t\}$, then for each given number $\alpha \in \mathbf{R}$ such that $\alpha < \sqrt{d}$, the set $\Sigma \cap (0, \alpha]$ is finite.

Here we give an example of a 3-dimensional integral hypersurface X of degree $d \geq 4$ in $\mathbf{P}^4$, which is smooth at a point 0 and it computes $\epsilon(\mathcal{O}_X(1), 0) = \frac{d}{d-1}$.

Example 3.1. *For any $d \geq 4$, there exists an integral hypersurface $X \subset \mathbf{P}^4$ containing no line through a point $0 \in X$ and smooth at 0 such that*

$$\epsilon(\mathcal{O}_X(1), 0) = \frac{d}{d-1}.$$

Construction of 3.1. The construction is a straight forward extension of Bauer's example [Bau99, Lemma 2.2].

We start with a surface S from the example 2.6. Let the defining equation of $S|_U$ is:

$$g(x_1, x_2, x_3) = 0$$

where $U \cong \mathbf{A}^3$ is the affine open neighborhood containing the point 0 . Choose a general smooth integral surface S' in $\mathbf{P}^3$ of degree $d-1$ satisfying

- (1) S' does not contain the point $0 = (0, 0, 0)$.
- (2) S' does not contain any 1-dimensional singular locus component of S .
- (3) If a point p is in the intersection of S and S' , and if S is smooth at p , then the tangent planes $T_p(S')$ and $T_p(S)$ are not equal.

Let the defining equation of S' be

$$g'(x_1, x_2, x_3) = 0.$$

Then, we define a hypersurface X in the affine open neighborhood $\mathbf{A}^4 \supset U$ by the equation

$$f = g + x_4 g' = 0.$$

We claim that X is smooth at the point $0 = (0, 0, 0, 0)$ and X is smooth except perhaps finitely many points.

Suppose X is singular at the point $p = (p_1, p_2, p_3, p_4)$. The partial derivative of f with respect to x_4 gives $g'(p_1, p_2, p_3) = 0$, and hence $f(p) = 0$ implies $g(p_1, p_2, p_3) = 0$. Therefore, the point $p' = (p_1, p_2, p_3)$ is an intersection point of S and S' . By the condition (1), X must be smooth at the point 0 .

Let $j_p^1(h)$ be the linear term of $\theta(h)$ where θ is a linear translation of the point 0 to the point $p = (p_1, p_2, p_3, p_4)$ where h is a polynomial in x_i for $1 \leq i \leq 4$.

Since X is singular at p , then $j_p^1(f) = 0$. On the other side,

$$0 = j_p^1(f) = j_{p'}^1(g) + p_4 \cdot j_{p'}^1(g').$$

If $p_4 \neq 0$ and $j_{p'}^1(g) = 0$, then

$$j_{p'}^1(g') = 0.$$

This is impossible when S' is smooth.

If $p_4 \neq 0$ and $j_{p'}^1(g) \neq 0$, then

$$T_{p'}(S') = T_{p'}(S).$$

This cannot be true by the condition (3).

If $p_4 = 0$, then $j_{p'}^1(g) = 0$. It means S is singular at p' . But only finitely many intersection points of S and S' can be in this situation by the condition (2). This completes the proof of the claim.

Since S is singular at 0, it is clear that the hyperplane $\{x_4 = 0\} = T_0X$ and hence $S = X \cap T_0X$.

Since X is singular at most finitely many points, then X is irreducible.

Moreover, X does not contain a line through 0. To this end, assume to the contrary that X contains a line through 0. Let the line be $\ell := \{(a_1t, a_2t, a_3t, a_4t) | t \in \mathbf{C}\}$. Then,

$$f(a_1t, a_2t, a_3t, a_4t) = 0 \quad \forall t.$$

However, if $a_4 \neq 0$, then it is impossible when g' has nonzero constant term, and g has no constant term and no linear term. If $a_4 = 0$, then the line $\ell \subset S$. This is not possible when S contains no line through 0. Therefore, we have

$$\epsilon(\mathcal{O}_X(1), 0) = \epsilon(\mathcal{O}_S(1), 0) = \frac{d}{d-1}.$$

□

REFERENCES

- [Bau99] Thomas Bauer. Seshadri constants on algebraic surfaces. *Math. Ann.*, 313(3):547–583, 1999.
- [Dem92] Jean-Pierre Demailly. Singular Hermitian metrics on positive line bundles. In *Complex algebraic varieties (Bayreuth, 1990)*, volume 1507 of *Lecture Notes in Math.*, pages 87–104. Springer, Berlin, 1992.
- [Laz04] Robert Lazarsfeld. *Positivity in algebraic geometry. I*, volume 48 of *[Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 2004. Classical setting: line bundles and linear series.
- [Ogu02] Keiji Oguiso. Seshadri constants in a family of surfaces. *Math. Ann.*, 323(4):625–631, 2002.